

# **VISUAL GUIDE**

# **Winter Wheat**

DEVELOPMENT AND GROWTH STAGING



University of Wisconsin-Madison | UW Extension

[WWW.COOLBEAN.INFO](http://WWW.COOLBEAN.INFO)

# Contents

Introduction

Basic wheat anatomy

Feekes scale for cereal growth stages

Zadoks scale for cereal growth stages

## ILLUSTRATED GROWTH STAGES

|       |                    |
|-------|--------------------|
| ..... | <b>Germination</b> |
| ..... | Feekes 1           |
| ..... | Feekes 2           |
| ..... | Feekes 3           |
| ..... | Feekes 4           |
| ..... | Feekes 5           |
| ..... | Feekes 6           |
| ..... | Feekes 7           |
| ..... | Feekes 8           |
| ..... | Feekes 9           |
| ..... | Feekes 10          |
| ..... | Feekes 10.1        |
| ..... | Feekes 10.2        |
| ..... | Feekes 10.3        |
| ..... | Feekes 10.4        |
| ..... | Feekes 10.5        |
| ..... | Feekes 10.5.1      |
| ..... | Feekes 10.5.2      |
| ..... | Feekes 10.5.3      |
| ..... | Feekes 10.5.4      |
| ..... | Feekes 11.1        |
| ..... | Feekes 11.2        |
| ..... | Feekes 11.3        |
| ..... | Feekes 11.4        |



### Other cereals



|       |                  |
|-------|------------------|
| ..... | <b>Barley</b>    |
| ..... | <b>Oats</b>      |
| ..... | <b>Rye</b>       |
| ..... | <b>Triticale</b> |

# Introduction

Understanding the growth stages of cereals crops and how to identify them is key to successful cropping and pest management decisions.

Although there are several growth staging methods, this guide is based on the Feekes scale, which is a popular tool used in the field. It has eleven development stages with some stages having more detailed subdivisions.

The Zadoks scale is the standard scale used in research and has ten development stages, each stage having ten subdivisions. Both scales are useful to know, so this guide cross-references the Zadoks equivalents to the Feekes.

***This guide uses winter wheat as an example.***

However, the methods generally apply to other cereals as well and at the back of the guide are sections that showcase barley, oats, rye and triticale.

## **A few notes on growth staging plants:**

- Select plants that represent at least 50% of the field
- Dig plants (if possible), so you can assess the entire plant
- Start at the base of plant and work your way upward
- Use a knife to split the stems and sheaths
- Look and feel for nodes

## **References:**

Large, E. C. (1954). *Growth Stages in Cereals Illustration of the Feekes Scale*. Plant Pathology, 3: 128–129. doi:10.1111/j.1365-3059.1954.tb00716.x



Feekes, Willem (1941). *De Tarwe en Haar Milieu [Wheat and Its Environment]*. Verslagen van de Technische Tarwe Commissie. (in Dutch (English summary)). 17: 523–888.

J.C. Zadoks, T.T. Chang, C.F. Konzani. *A Decimal Code for the Growth Stages of Cereals*. Weed Research 1974 14:415-421.

# Basic wheat anatomy

This illustration shows the basic anatomy of a Feekes 2 plant; the intention of the drawing is to illustrate and annotate an ideal plant

Plants in the field are rarely ideal and are often missing leaves, tillers, etc.



During the vegetative stages, the **auricles** are often a good way to distinguish between different cereal crops

## Wheat

Auricles blunt and hairy; leaf sheath and blade always hairy; ligule medium length; leaf blades twist clockwise

# Feekes scale for cereal growth stages

|                 |        |                                                                                                 |
|-----------------|--------|-------------------------------------------------------------------------------------------------|
| SEEDLING GROWTH | 1      | One shoot, first leaf through coleoptile                                                        |
| TILLERING       | 2      | Tillering begins; main shoot and one tiller                                                     |
|                 | 3      | Tillers formed; leaves often twisted<br>In some varieties, plant may be prostrate in appearance |
|                 | 4      | Leaf sheaths lengthen; beginning pseudostem erection                                            |
|                 | 5      | Leaf sheaths fully elongated to form strongly erect pseudostem                                  |
| STEM EXTENSION  | 6      | First node of stem visible at base of shoot; jointing                                           |
|                 | 7      | Second node of stem formed; next-to-last leaf just visible                                      |
|                 | 8      | Flag leaf visible but still rolled up                                                           |
|                 | 9      | Ligule of flag leaf just visible                                                                |
|                 | 10     | Flag leaf sheath completely grown out; booting                                                  |
| HEADING         | 10.1   | First awns of head just visible                                                                 |
|                 | 10.2   | 1/4 of heading process complete                                                                 |
|                 | 10.3   | 1/2 of heading process complete                                                                 |
|                 | 10.4   | 3/4 of heading process complete                                                                 |
|                 | 10.5   | All heads out of sheath                                                                         |
| FLOWERING       | 10.5.1 | Beginning of flowering                                                                          |
|                 | 10.5.2 | Flowering complete to top of head                                                               |
|                 | 10.5.3 | Flowering complete at base of head                                                              |
|                 | 10.5.4 | Flowering complete; kernel watery ripe                                                          |
| RIPENING        | 11.1   | Kernel milky ripe; milk stage                                                                   |
|                 | 11.2   | Kernel mealy ripe; soft but dry consistency; soft dough stage                                   |
|                 | 11.3   | Kernel hard; difficult to divide with thumbnail; hard dough stage                               |
|                 | 11.4   | Kernel harvest ready; straw dead                                                                |

# Zadoks scale

for cereal growth stages

|                 |    |                                      |                               |     |
|-----------------|----|--------------------------------------|-------------------------------|-----|
| Germination     | 00 | Dry seed                             | Feekes<br>scale<br>equivalent |     |
|                 | 01 | Start of imbibition                  |                               |     |
|                 | 03 | Imbibition complete                  |                               |     |
|                 | 05 | Radicle emerged from seed            |                               |     |
|                 | 07 | Coleoptile emerged from seed         |                               |     |
|                 | 09 | Leaf just at coleoptile tip          |                               |     |
| Seedling growth | 10 | First leaf through coleoptile        | TILLERING                     | 1   |
|                 | 11 | First leaf unfolded                  |                               |     |
|                 | 12 | 2 leaves unfolded                    |                               |     |
|                 | 13 | 3 leaves unfolded                    |                               |     |
|                 | 14 | 4 leaves unfolded                    |                               |     |
|                 | 15 | 5 leaves unfolded                    |                               |     |
|                 | 16 | 6 leaves unfolded                    |                               |     |
|                 | 17 | 7 leaves unfolded                    |                               |     |
|                 | 18 | 8 leaves unfolded                    |                               |     |
|                 | 19 | 9 or more leaves unfolded            |                               |     |
| Tillering       | 20 | Main shoot only                      | TILLERING                     | 2   |
|                 | 21 | Main shoot and 1 tiller              |                               |     |
|                 | 22 | Main shoot and 2 tillers             |                               |     |
|                 | 23 | Main shoot and 3 tillers             |                               |     |
|                 | 24 | Main shoot and 4 tillers             |                               | 3   |
|                 | 25 | Main shoot and 5 tillers             |                               |     |
|                 | 26 | Main shoot and 6 tillers             |                               |     |
|                 | 27 | Main shoot and 7 tillers             |                               |     |
|                 | 28 | Main shoot and 8 tillers             |                               |     |
|                 | 29 | Main shoot and 9 or more tillers     |                               |     |
| Stem elongation | 30 | Pseudostem erection                  | STEM EXTENSION                | 4-5 |
|                 | 31 | 1 <sup>st</sup> node detectable      |                               | 6   |
|                 | 32 | 2 <sup>nd</sup> node detectable      |                               | 7   |
|                 | 33 | 3 <sup>rd</sup> node detectable      |                               |     |
|                 | 34 | 4 <sup>th</sup> node detectable      |                               |     |
|                 | 35 | 5 <sup>th</sup> node detectable      |                               |     |
|                 | 36 | 6 <sup>th</sup> node detectable      |                               |     |
|                 | 37 | Flag leaf just visible               |                               | 8   |
|                 | 39 | Flag leaf ligule/collar just visible |                               | 9   |

A leaf is unfolded when its ligule is visible, or the tip of the next leaf is visible

|                         |    |                                                  |                |        |
|-------------------------|----|--------------------------------------------------|----------------|--------|
| Booting                 | 40 | -----                                            | STEM EXTENSION | 10     |
|                         | 41 | Flag leaf sheath extending                       |                |        |
|                         | 45 | Boot just visibly swollen                        |                |        |
|                         | 47 | Flag leaf sheath opening                         |                |        |
|                         | 49 | First awns visible                               |                |        |
| Inflorescence emergence | 50 | First spikelet of inflorescence visible          | HEADING        | 10.1   |
|                         | 53 | 1/4 of inflorescence emerged                     |                | 10.2   |
|                         | 55 | 1/2 of inflorescence emerged                     |                | 10.3   |
|                         | 57 | 3/4 of inflorescence emerged                     |                | 10.4   |
|                         | 59 | Emergence of inflorescence completed             |                | 10.5   |
| Anthesis                | 60 | Beginning of anthesis                            | FLOWERING      | 10.5.1 |
|                         | 65 | Anthesis half-way                                |                | 10.5.2 |
|                         | 69 | Anthesis completed                               |                | 10.5.3 |
| Milk development        | 70 | -----                                            |                |        |
|                         | 71 | Kernel watery ripe                               |                | 10.5.4 |
|                         | 73 | Early milk                                       |                |        |
|                         | 75 | Medium milk                                      | RIPENING       | 11.1   |
|                         | 77 | Late milk                                        |                |        |
| Dough development       | 80 | -----                                            |                |        |
|                         | 83 | Early dough                                      |                |        |
|                         | 85 | Soft dough                                       |                | 11.2   |
|                         | 87 | Hard dough                                       |                |        |
|                         | 90 | -----                                            |                |        |
| Ripening                | 91 | Kernel hard (difficult to divide with thumbnail) |                | 11.3   |
|                         | 92 | Kernel hard (no longer dented with thumbnail)    |                | 11.4   |
|                         | 93 | Kernel loosening in daytime                      |                |        |
|                         | 94 | Overripe, straw dead and collapsing              |                |        |
|                         | 95 | Seed dormant                                     |                |        |
|                         | 96 | Viable seed giving 50% germination               |                |        |
|                         | 97 | Seed not dormant                                 |                |        |
|                         | 98 | Secondary dormancy induced                       |                |        |
|                         | 99 | Secondary dormancy lost                          |                |        |

## Germination

Germination begins when the **dry seed** imbibes water and begins to expand

**Zadoks 00**  
Dry seed



**Zadoks 01**  
Start of imbibition



**radicle**  
The first root



**Zadoks 05**  
Radicle emerged from seed


**coleoptile**  
The round sheath that protects the first leaf



**Zadoks 07**  
Coleoptile emerged from seed

**Planting depth and soil temperature**  
influence the duration of the  
germination stages

Zadoks  
00-09



# Germination

The **first true leaf** emerges through the coleoptile's tip

**Zadoks 09**  
Leaf just at coleoptile tip

The **coleoptile** stops growth when it encounters light above the soil surface

**seed**

The **seminal roots** begin developing

Seminal roots are fibrous and are produced by the seed



A close-up photograph of several young corn plants. The plants are green and have one leaf visible through the coleoptile. They are growing out of dark, moist soil. The background is a bright blue sky with some light clouds.

Feekes

1

Zadoks 10

EMERGENCE | **Feekes 1** | One shoot formed;  
first leaf through coleoptile

Feekes  
1

This is an important time to check plants for **uniform emergence**; planting depth and soil temperature influence the length of this stage

main shoot





Feekes  
2

Zadoks 21

TILLERING | **Feekes 2** | Tiller development begins

Tillers produced  
**in the fall** will  
contribute more  
to grain yield than  
those produced in  
the spring

Feekes  
**2**

main shoot

tillers

The **crown** forms  
between the seed  
and soil surface



Feekes

2

main shoot

tiller

tillers

**prophyll**

The independent  
sheath at the  
base of each tiller

tiller

The secondary  
root system starts  
developing



# Are tillers important?

**Tillers are absolutely necessary for high yields**

Feekes  
2

**1** can  
produce  
**planted seed**

**4-5**

**tillers**

are also called  
axillary or  
side shoots;  
not all tillers  
will complete  
development  
and produce  
grain

The **total numbers of tillers**  
a plant produces is determined by both  
**environmental conditions** and **genetic potential**

## KEY YIELD COMPONENT

A **tiller** is capable of forming a single head (spike)

The **head** is made up of spikelets

Each **spikelet** contains individual florets

Individual **florets** can produce a single **kernel**



In Wisconsin, the recommended  
planting date range for optimal  
tiller development in winter wheat  
is **September 20 to October 10**

Feekes  
3



Zadoks  
22-29

TILLERING | Feekes 3 | Tillering completed


Tillers with 3 or more leaves will be nutritionally independent from the main shoot

Feekes  
3

main shoot

★ tillers

The secondary root system undergoes extensive development



Feekes  
3

## FEEKES 3 CAN OCCUR IN FALL OR SPRING

because winter wheat development is dependent on both temperature and planting date

### FEEKES 3 FALL

Tillering completes in fall,  
winter dormancy occurs

### FEEKES 3 SPRING

Tillering begins in the fall,  
winter dormancy occurs,  
tillering completes in the spring

*What happens during  
winter dormancy?*

# Vernalization!

When temperatures fall below 50 degrees Fahrenheit  
for typically 3-6 weeks, the plant initiates

## differentiation



**the growing point changes** from  
vegetative or leaf producing to  
reproductive or spikelet producing



*The growing point is at the **double ridge stage** and  
is still protected in the crown below the soil surface*

*The number of florets initiated during this stage will  
determine the **potential** number of kernels per head*

**KEY YIELD COMPONENT**



Dig plants as soon as the soil thaws, bring inside and place in a warm (preferably moist) area for a few days, then check for root regrowth



#### regrowth

Root regrowth will develop from the crown and appear a vibrant white compared to the older roots

Feekes

3



## FACTORS AFFECTING WINTER SURVIVAL

-  Good snow cover acts as insulator; keeps soil temperature from going below critical levels
-  Cyclic freezing and thawing increases injury from ice crystal growth in tissue
-  Mid-winter thaw and rain cause flooding at the base of the plants; crowns can die at warmer temperatures
-  Ice encasement traps carbon dioxide and suffocates plant by inhibiting respiration
-  Frost heaving can push root system out of ground, leaving plants vulnerable and weak

# 4 STEPS TO ASSESS STANDS IN EARLY SPRING

- 1 Venture out and get a general overview of the fields** — vibrant green patches may be interspersed with drab brown areas, but brown does not always indicate winter-killed plants
- 2 Check for winter survival** — identify several representative plants and 1) dig plants and bring inside to check for root regrowth or 2) wait a week and revisit to check for regrowth in the field
- 3 Do a plant count** — below 12 live plants per square foot is an automatic replant; 12-15 live plants per square foot requires more consideration for a replant decision; 15-22 live plants per square foot may recover and reach maximum yield potential; over 22 live plants per square foot means you're good to go!
- 4 Consider a nitrogen application** — the optimal time to apply nitrogen to winter wheat in Wisconsin is during green up; for recommendations and rates, consult UW-Extension publication A2809 *Nutrient Application Guidelines for Field, Vegetable and Fruit Crops in Wisconsin*

In Wisconsin, the growth stage at green up can be **Feekes 3** or **Feekes 4** depending on planting date and environmental conditions

**GREEN  
UP**

### → **How to do a plant count**

Count the number of plants in a 3-foot length

Do this for at least 3 areas

Take the average of the counts

Multiply that number by 4

Then divide by the row width (inches)

#### **EXAMPLE**

*The 3 counted areas have  
40, 35 and 45 plants*

*Add 40, 35 and 45 and then  
divide by 3, the average = 40*

*Multiply  $40 \times 4 = 160$*

*Divide 160 by 7.5 inches =  
**21 plants/square foot***



Feekes

4

Zadoks 30

TILLERING | **Feekes 4** | Leaf sheaths lengthen,  
pseudostem erection begins

This is an important time for weed control  
and/or nitrogen applications

Wheat plants have a *pseudostem*, which is a false  
stem composed of concentric rolled leaf sheaths  
that surround the growing point (the developing head)

During this stage, these leaf sheaths lengthen,  
making the plants stand more upright

Feekes

4



This is the last stage that some herbicides  
can be used without risk of injury!

*Always check and follow herbicide labels*

Feekes

5



Zadoks 30


TILLERING | **Feekes 5** | Leaf sheaths fully  
elongated, pseudostem strongly erect



Feekes  
5

plants cut at  
soil surface

As the **developing head** is pushed up into the pseudostem, it becomes more vulnerable to damage



less than 1/8 inch

The growing point is at the **terminal spikelet stage** and about 1/4 inch above the crown

The **number of spikelets per head** has been determined by this stage

KEY YIELD COMPONENT

Feekes  
6

1<sup>st</sup> node

1<sup>st</sup> node

*As the head moves up the stem, it is vulnerable to freeze injury during low temperatures!*

## Feekes 6

1<sup>st</sup> node




**A node** is an area of active cell division from which leaves, tillers and adventitious roots develop

1<sup>st</sup> node




leaves  
removed to  
show nodes



The jointing stage is when the **internodal tissue** begins to elongate and pushes the **four nodes** that are stacked in the crown upward, similar to how a telescope works

A leaf arises from each of these nodes, with the 4<sup>th</sup> node giving rise to the flag leaf — the last leaf the plant produces



Feekes 6 plant  
shown with all  
leaves intact



Feekes  
6

*From this growth stage forward,  
**broken stems due to wheel  
traffic will result in yield loss!***

*The developing head is  
moving up the stem and  
needs to be protected*

***The number of tillers that form heads  
has been determined by this stage***

**KEY YIELD COMPONENT**

Feekes  
7

## **SCOUT NOW!**

THIS WILL GIVE YOU  
THE INFORMATION  
YOU NEED TO MAKE  
GOOD MANAGEMENT  
DECISIONS ABOUT  
PROTECTING THE  
FLAG LEAF AT THE  
NEXT STAGE

1<sup>st</sup> node

2<sup>nd</sup> node

STEM EXTENSION | Feekes 7 | Two nodes  
visible above the soil line

## Feekes 7

This leaf arises  
from the 2<sup>nd</sup> node

location of  
developing  
head

This leaf arises  
from the 1<sup>st</sup> node

To demonstrate this, pull  
the leaf sheath back and  
downward; it will break off  
at the node

2<sup>nd</sup>  
node

4<sup>th</sup>  
node

3<sup>rd</sup>  
node

sheaths removed and  
stem slit to show head  
(about 1-1/2 inches)  
and nodes

1<sup>st</sup> node



Feekes  
8

flag leaf

This is a **critical time** to make foliar fungicide application decisions!

flag leaf



### FLAG LEAF FACTS

The flag leaf accounts for over 50% of the photosynthates used for grain development, a.k.a **YIELD**

It must be protected from disease or insect damage to ensure the plant's full yield potential

Fungicide application decisions to protect the flag leaf should be made based on **presence** and **severity** of disease on the two leaves immediately below it

Feekes  
8

The **flag leaf** arises from the 4<sup>th</sup> node

#4

This leaf arises from the 2<sup>nd</sup> node

#2

This leaf arises from the 3<sup>rd</sup> node

#3

### HOW DO YOU KNOW IF IT'S THE FLAG LEAF?

Identify the leaf arising from the 1<sup>st</sup> node

Call this leaf #1 and count upward

The flag leaf will be leaf #4

location of developing head

4<sup>th</sup> node

3<sup>rd</sup> node

2<sup>nd</sup> node

#1

This leaf arises from the 1<sup>st</sup> node  
(see tip on Feekes 7)

sheaths removed to show head and nodes

**Kernel size** is determined by crop health and water/nutrient availability beginning now and continuing through grain fill

KEY YIELD COMPONENT


Feekes

9



Zadoks 39

STEM EXTENSION | **Feekes 9** | Flag leaf fully emerged from the whorl; ligule just visible



**flag leaf**

**ligule**

**leaf collar**

The area on the outer side of the leaf where the blade and the sheath join

**The ligule** is a narrow membranous scale on the inner side of the leaf sheath at its junction with the blade

Feekes  
9

***CONTINUE TO SCOUT FOR  
INSECT PESTS AND DISEASES !***

Feekes  
**10**

**CONTINUE TO SCOUT  
FOR INSECT PESTS  
AND DISEASES !**

location of  
developing  
head

**At this stage, the Feekes  
scale subdivides:**

- 10.1 Head emerging
- 10.2 Heading 1/4 complete
- 10.3 Heading 1/2 complete
- 10.4 Heading 3/4 complete
- 10.5 Heading complete

**and then subdivides again**

- 10.5.1 Beginning flowering
- 10.5.2 Flowering complete to top of spike
- 10.5.3 Flowering complete at base of spike
- 10.5.4 Kernels watery ripe

STEM EXTENSION | **Feekes 10** | Flag leaf sheath  
completely grown out; head visible in the  
leaf sheath; booting

Feekes  
10

sheath  
removed to  
show detail  
of developing  
head

spikelets


awns

approximately 3 inches long

peduncle

The stem  
that  
supports  
the head

The developing  
head is pushed  
through the flag  
leaf sheath as the  
peduncle and  
sheath elongate



Feekes  
10.1

**Awns** are the slender bristles that extend from the floret; some wheat varieties are awnless (also called beardless)

As the leaf sheath splits, the awns become visible

During head emergence, the tiller's development synchronizes with the main stem

The result is that flowering occurs simultaneously throughout the plant, even though the tillers may have emerged at different times

HEADING | **Feekes 10.1** | First awns of the head just visible

When determining the growth stage of a field, **50% of the plants** must be at that stage or above

Feekes  
10.1

Feekes  
10.1

Feekes  
10.2





Feekes  
10.2

Zadoks 53

HEADING | **Feekes 10.2** | 1/4 of the head  
emerged from the leaf sheath



Feekes  
10.1

Feekes  
10.2


Feekes  
10.2

sheaths removed  
to show  
developing  
heads

Feekes  
10.3



HEADING | **Feekes 10.3** | 1/2 of the head  
emerged from the leaf sheath

Zadoks 55



Feekes  
10.3

### spikelet opened to show detail





Feekes  
10.4

Zadoks 57

HEADING | **Feekes 10.4** | 3/4 of the head  
emerged from the leaf sheath

Feekes  
10.4



head lifted out of sheath to  
show elongating peduncle

Feekes  
10.5



Zadoks 59

HEADING | **Feekes 10.5** | Head completely  
emerged from the leaf sheath

The image shows three wheat spikes against a white background. The spikes are green and elongated, with the central spike being the most prominent. The glumes are tightly packed, and the awns are visible at the tips. The spikes are shown from a slightly low angle, giving them a sense of height and growth.

This stage completes the heading process

Feekes  
10.5

Feekes

10.5.1

*Starting now and continuing 5-7 days **after this stage** is the optimum time for fungicide application to suppress *Fusarium* head blight (FHB), also called head scab*



*Flowering begins slightly above the middle portion of the head and continues towards the top*

This stage begins shortly after the head has completely emerged from the leaf sheath

Anthers dangle outside the florets during flowering

awn

Feekes

10.5.1

*The number of **flowers pollinated** determines the number of **kernels** that will develop*

#### KEY YIELD COMPONENT

##### **anther**

The male flower part that produces and releases pollen

##### **pollen**

The powder-like grains that enable fertilization

Feekes

10.5.2

The *developing head* is still vulnerable to freeze injury during low temperatures

Flowering continues toward the base

FLOWERING | Feekes 10.5.2 | Flowering complete to the top of the head



**Anthers** fade to white as flowering completes at the top of the head, while those toward the base are still brightly colored

Feekes

10.5.2

floret opened shortly after pollination to show developing kernel

Feekes

10.5.3



Zadoks 69

FLOWERING | **Feekes 10.5.3** | Flowering  
complete at the base of the head



This stage  
signals  
the end of  
pollination

Feekes  
10.5.3



floret outer structure removed to  
show developing kernel

This is the beginning of the **grain filling stages**;  
kernel length is established during this stage

Feekes

10.5.4





Feekes

10.5.4

When squeezed, *clear fluid*  
is released from the kernel



developing kernel with  
desiccated anthers  
still attached



*Kernel size increases*

but not dry matter  
accumulation



Feekes

11.1

Zadoks 75

RIPENING | **Feekes 11.1** | Kernel milky ripe;  
milk stage

Feekes

11.1



Dry matter accumulates  
in the kernel

When squeezed,  
**milk-like fluid**  
is released from  
the kernel





Feekes  
11.2

Zadoks 85

RIPENING | Feekes 11.2 | Kernel mealy ripe;  
soft but dry consistency; soft dough stage



Feekes

11.2


Green color of the kernel, glume and peduncle begins to fade

***Starch, nutrients and dry matter***  
accumulate rapidly in the kernel



The kernel's content is a soft-doughy material





Feekes  
11.3

Zadoks 91

RIPENING | **Feekes 11.3** | Kernel hard; difficult to divide with a thumbnail; hard dough stage



Feekes  
11.3

Kernels reach their **maximum dry weight** and are **physiologically mature**



Kernel  
moisture  
decreases from  
40% to 30%



Feekes

11.4



Kernel  
moisture  
decreases from  
30% to 15%



Zadoks 92

RIPENING | **Feekes 11.4** | Kernel harvest ready;  
straw dead

Green plant tissue fades to *straw*

Feekes

11.4





Other  
cereals



# Barley

Auricles long, slender and hairless; leaf sheath and blade usually hairless (scattered hairs on some varieties); ligule medium length; leaf blades twist clockwise



Other  
cereals

Other  
cereals



*To distinguish oats  
from wheat during the  
vegetative stages, check  
the auricles — oats lack  
auricles, while wheat  
auricles are blunt and hairy*



# Oats

Auricles absent; leaf sheath and blade hairless (scattered hair on some varieties); ligule medium length; leaf blades twist counter-clockwise

The background of the entire page is a close-up photograph of several oat panicles. The panicles are light brown and tan, with long, thin awns extending from them. The leaves are green and have prominent parallel veins. The overall composition is vertical and artistic, with a soft, natural lighting.

Other  
cereals



Oats

*A simple method to distinguish oats from all other cereals during the vegetative stages is to observe the twist of the leaves; when viewing from above, oat leaves will have a counter-clockwise curl, all other cereals' leaves curl clockwise*



All other cereals



Other  
cereals



*To distinguish rye  
from wheat during the  
vegetative stages, check  
the auricles — rye auricles  
are short and hairless,  
while wheat auricles are  
blunt and hairy*



# Rye

Auricles very short and hairless; leaf sheath and blade have an inconsistent degree of hairiness; ligule short; leaf blades twist clockwise



Other  
cereals



Other  
cereals




*The auricles of both triticale and wheat are blunt and hairy, so they are difficult to distinguish from each other during the vegetative stage*

*An alternative method is to remove a seedling from the soil and check the grain shell; triticale shells are oblong in shape and dark in color, while wheat grain shells are oval and lighter*



# Triticale

Auricles blunt and hairy, leaf sheath and blade hairy; ligule of medium length; leaf blades twist clockwise



Other  
cereals

## Authors

Mimi Broeske, Nutrient and Pest Management Program; Shawn Conley, John Gaska and Adam Roth, Department of Agronomy; University of Wisconsin-Madison, College of Agricultural and Life Sciences and Division of Extension

## Acknowledgements

The authors would like to thank our external reviewers:

Mike Cerny, Walworth County farmer

Dr. Chad Lee, University of Kentucky

Dr. David Marburger, Oklahoma State University



This publication is available from the  
**Nutrient and Pest Management Program:**  
web ([ipcm.wisc.edu](http://ipcm.wisc.edu)); phone (608) 265-2660;  
email ([npm@hort.wisc.edu](mailto:npm@hort.wisc.edu))

An EEO/AA employer, University of Wisconsin-Madison Division of Extension provides equal opportunities in employment and programming, including Title VI, Title IX, the Americans with Disabilities Act (ADA) and Section 504 of the Rehabilitation Act requirements.

